



**Cancer Epidemiology Services
Public Health Services Branch
New Jersey Department of Health and Senior Services**

**Probability of Developing Cancer for Selected Age Groups by Sex,
2001-2003 (New Jersey and U.S.)**

Technical Notes

The following table presents probabilities (risk estimates) of developing invasive cancers from birth to death (all ages) and for the age intervals 0-39, 40-59 and 60-79 for New Jersey and the U.S. These probabilities are based on cancers diagnosed from 2001-2003. The cancer risk estimates are presented for men and women separately for all sites combined and for fourteen common cancer subtypes. The data represent all races combined with the exception of melanoma of the skin which is presented for whites only. In addition, *in situ* cancers are not included except for bladder cancer *in situ* cases, which are included with invasive urinary bladder and all sites.

The probabilities of developing invasive cancers for selected age intervals were calculated using DevCan Software¹. DevCan calculates the probability of developing or dying of cancer using both incidence and mortality data².

New Jersey cancer incidence data for first primary cancers only were taken from the November 2005 analytic file of the New Jersey State Cancer Registry. New Jersey cancer mortality data were obtained through the SEER Program website. With the exception of the two additional criteria used for this analysis (year of diagnosis 2001-2003 and first primary cancers only), the data specifications for New Jersey incidence and mortality data used for this analysis were the same as the specifications in our recent report *Cancer Incidence and Mortality in New Jersey, 1999-2003*³. The New Jersey incidence and mortality data for 2001-2003 were imported into DevCan to create a New Jersey DevCan database for calculating the probabilities of developing cancer.

The U.S. DevCan database was downloaded directly from National Cancer Institute (NCI) Surveillance, Epidemiology, and End Results (SEER) Program⁴.

How to Read the Table

The probability of developing invasive cancer from birth to death, also known as lifetime risk, among New Jersey residents is about 60% for men and 49% for women, both of which round off to 1 in 2. The probability of developing invasive colorectal cancer between the ages 40-59, among New Jersey residents is 1.02% for men and 0.80% for women, which round off to 1 in 98 men and 1 in 125 women.

These probabilities reflect the average experience of people in New Jersey and do not take into account individual behaviors and risk factors. For example, the estimate of 1 man in 13 developing lung cancer in a lifetime underestimates the risk of smokers and overestimates the risk for nonsmokers⁵.

For additional information contact Pamela Agovino or Xiaoling Niu of the Cancer Epidemiology Services, New Jersey Department of Health and Senior Services. They can be reached at 609-588-3500 or by email (Pamela.Agovino@doh.state.nj.us, Xiaoling.Niu@doh.state.nj.us).

Probability of Developing Invasive Cancers for Selected Age Groups by Sex, 2001-2003 (New Jersey and U.S.)*

		New Jersey								U.S.							
		All Ages**		0-39		40-59		60-79		All Ages **		0-39		40-59		60-79	
	Sex	%	1 in	%	1 in	%	1 in	%	1 in	%	1 in	%	1 in	%	1 in	%	1 in
All Sites ***	Male	60.04	2	1.68	60	10.40	10	46.26	2	45.51	2	1.42	71	8.50	12	34.30	3
	Female	48.55	2	2.39	42	11.02	9	29.40	3	38.01	3	2.02	49	8.98	11	22.29	4
Oral Cavity and Pharynx	Male	1.29	77	0.04	2,458	0.42	239	0.76	132	1.38	73	0.04	2,390	0.41	241	0.83	120
	Female	0.63	160	0.03	3,001	0.14	725	0.35	288	0.67	149	0.03	3,006	0.15	686	0.36	280
Colon and Rectum	Male	6.68	15	0.08	1,274	1.02	98	4.46	22	5.75	17	0.08	1,339	0.91	111	3.82	26
	Female	6.29	16	0.09	1,169	0.80	125	3.43	29	5.42	18	0.07	1,522	0.70	142	2.92	34
Lung and Bronchus	Male	8.08	12	0.03	2,944	1.13	89	5.99	17	7.62	13	0.03	3,293	0.97	103	5.69	18
	Female	6.48	15	0.05	2,031	0.98	102	4.34	23	5.82	17	0.03	3,216	0.78	128	3.96	25
Melanoma of the Skin (White)	Male	2.54	39	0.15	667	0.64	156	1.52	66	2.33	43	0.15	663	0.61	165	1.34	75
	Female	1.74	58	0.27	374	0.49	204	0.79	127	1.63	61	0.27	365	0.51	198	0.68	148
Breast	Female	13.04	8	0.56	180	4.24	24	7.06	14	12.82	8	0.48	209	3.96	25	7.16	14
Cervix Uteri	Female	0.81	124	0.16	623	0.33	304	0.28	359	0.72	140	0.15	665	0.28	364	0.25	402
Corpus and Uterus, NOS	Female	2.99	33	0.05	1,942	0.83	121	1.88	53	2.59	39	0.06	1,598	0.72	138	1.54	65
Ovary	Female	1.58	63	0.08	1,265	0.44	230	0.82	122	1.47	68	0.08	1,296	0.37	269	0.79	127
Prostate	Male	19.99	5	0.02	6,649	3.19	31	16.64	6	17.70	6	0.01	10,287	2.65	38	14.54	7
Urinary Bladder ***	Male	4.31	23	0.03	3,732	0.53	190	2.91	34	3.62	28	0.02	4,397	0.39	256	2.42	41
	Female	1.44	70	0.02	6,558	0.18	558	0.82	122	1.14	88	0.01	10,297	0.12	822	0.65	155
Brain and Other Nervous System	Male	0.71	142	0.14	734	0.18	555	0.37	270	0.66	153	0.12	822	0.16	639	0.35	290
	Female	0.56	179	0.12	861	0.12	806	0.26	384	0.52	193	0.11	900	0.11	918	0.24	420
Thyroid	Male	0.44	228	0.08	1,257	0.16	629	0.21	478	0.37	268	0.06	1,597	0.13	752	0.18	550
	Female	1.31	77	0.37	267	0.52	193	0.41	246	1.04	96	0.30	339	0.40	247	0.32	310
Non-Hodgkin Lymphoma	Male	2.38	42	0.16	641	0.50	202	1.38	72	2.23	45	0.14	736	0.46	218	1.34	75
	Female	1.97	51	0.10	1,002	0.36	275	1.09	92	1.88	53	0.09	1,138	0.32	311	1.04	97
Leukemia	Male	1.55	65	0.17	595	0.25	397	0.89	112	1.54	65	0.16	646	0.22	452	0.88	113
	Female	1.08	93	0.12	819	0.16	636	0.52	193	1.08	93	0.12	834	0.14	701	0.50	198

* For those free of cancer at beginning of age interval. Based on New Jersey and U.S. invasive cancer cases diagnosed during 2001-2003. Includes all races except melanoma of skin includes whites only.

** All Ages represents birth to death.

*** All Sites exclude basal and squamous cell skin cancers and *in situ* cancers except urinary bladder.

Probabilities of developing cancer were calculated using DevCan (Probability of Developing Cancer Software) developed by National Cancer Institute (NCI).

Data sources: New Jersey incidence data - New Jersey State Cancer Registry, New Jersey Department of Health & Senior Services 2005; New Jersey mortality data and U.S. incidence and mortality data, - Surveillance, Epidemiology, and End Results (SEER) Program, NCI; underlying mortality data - National Center for Health Statistics

References

1. DevCan: Probability of Developing or Dying of Cancer Software, Version 6.0.1. Statistical Research and Applications Branch, National Cancer Institute, 2005. [URL. <http://srab.cancer.gov/devcan>, accessed June 22, 2006]
2. Fay, M.P., Pfeiffer, R., Cronin, K.A., Le, C. and Feuer, E.J. (2003) "*Age-conditional probabilities of developing cancer*" *Statistics in Medicine*, 22(11): 1837-1848. Fay M.P. "*Estimating age conditional probability of developing disease from surveillance data*". *Population Health Metrics* 2004, 2:6. [URL. <http://www.pophealthmetrics.com/content/2/1/6>, accessed June 22, 2006]
3. Burger, S.S., Niu, X. et al *Cancer Incidence and Mortality in New Jersey, 1999-2003*. New Jersey Department of Health and Senior Services, June 2005. [URL. <http://nj.gov/health/ces/documents/report99-03.pdf> , accessed June 22, 2006]
4. Surveillance, Epidemiology, and End Results (SEER) Program (<http://www.seer.cancer.gov/>) DevCan database: "*SEER 13 Incidence and Mortality, 2000-2002, Follow-back Year-1992, with Kaposi Sarcoma and Mesothelioma*". National Cancer Institute, DCCPS, Surveillance Research Program, Cancer Statistics Branch, released November 2005, based on the November 2004 submission. Underlying mortality data provided by NCHS. [URL. <http://www.cdc.gov/nchs/>), accessed June 22, 2006]
5. American Cancer Society. *Cancer Facts & Figures 2006*. Atlanta: American Cancer Society; 2006. [URL. http://www.cancer.org/docroot/STT/content/STT_1x_Cancer_Facts_Figures_2006.asp, accessed June 22, 2006]